

N69-19186
CR-100255



APPROXIMATE TRANSITION PROBABILITIES AND
LIFETIME OF SOME OF THE EXCITED
STATES OF NEUTRAL IODINE

S.H. Koozekanani and G.L. Trusty

The Ohio State University
ElectroScience Laboratory
(formerly Antenna Laboratory)
Department of Electrical Engineering
Columbus, Ohio 43212

**CASE FILE
COPY**

TECHNICAL REPORT 1093-43
11 December 1968

Grant Number NGR-36-008-002

National Aeronautics and Space Administration
Office of Grants and Research Contracts
Washington, D.C. 20546

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The Government has the right to reproduce, use, and distribute this report for governmental purposes in accordance with the contract under which the report was produced. To protect the proprietary interests of the contractor and to avoid jeopardy of its obligations to the Government, the report may not be released for non-governmental use such as might constitute general publication without the express prior consent of The Ohio State University Research Foundation.

REPORT
by
The Ohio State University ElectroScience Laboratory
(Formerly Antenna Laboratory)
Columbus, Ohio 43212

| | |
|-------------------|--|
| Sponsor | National Aeronautics and Space Administration Office of Grants and Research Contracts Washington, D.C. 20546 |
| Grant Number | NGR-36-008-002 |
| Investigation of | Receiver Techniques and Detectors for Use at Millimeter and Submillimeter Wave Lengths |
| Subject of Report | Approximate Transition Probabilities and Lifetime of Some of the Excited States of Neutral Iodine |
| Submitted by | S.H. Koozekanani and G.L. Trusty ElectroScience Laboratory Department of Electrical Engineering |
| Date | 11 December 1968 |

ABSTRACT

Approximate transition probabilities and lifetime of some of the 5d, 6p, and 6s states of neutral iodine is given with the assumption that the states are purely pair coupled.

APPROXIMATE TRANSITION PROBABILITIES AND LIFETIME OF SOME OF THE EXCITED STATES OF NEUTRAL IODINE

Approximate transition probabilities for some of the excited levels of neutral iodine have been calculated. These data are useful in the approximate determination of line strength and prediction of possible new laser lines. The states are assumed to be pair coupled¹ and transition probabilities for $5p^4-5d \rightarrow 5p^4-6p$, $5p^4-6p \rightarrow 5p^4-6s$ and $5p^4-6s \rightarrow 5p^5$ configurations are given in Tables Ia through Ic, based upon the assumption that the states are not mixed. Table II gives the radials integrals

$$\int r \mathcal{F}_{nl}(r) \mathcal{F}_{nl'}(r) dr$$

which have been used in the calculations of the transition probabilities. Table III gives the exchange, direct and spin orbit interaction integrals for the three configurations $5p^4-5d$, $5p^4-6p$ and $5p^4-6s$ of iodine I. The above radial integrals were calculated using wavefunctions obtained from a computer program given by Herman and Skilman² for the solution of the self-consistent Hartree Fock equation in the Slater approximation. Finally in Table IV we have the lifetimes of some of the upper states.

In the calculations of the transition probabilities from the $5p^4-6s$ and $5p^4-5d$ states to the ground $5p^5 \ ^2P_J$ state one has first to express the ground $|5p^5 \ ^2P_J\rangle$ state in terms of its parents, i.e.,

$$|\ell^5 LSJM\rangle = \sum_{\bar{L} \bar{S}} (\ell^5 LS \mid \ell^4 \bar{L} \bar{S}) \{ \mid \ell^4 \bar{L} \bar{S}; \ell_s, LSJM \rangle$$

or

$$= \sum_{\bar{L} \bar{S}} \sum_{\bar{J} K} (-1)^{2\bar{S} + \bar{L} + \ell + \frac{1}{2} + K + L + J} (\ell^5 LS \{ \mid \ell^4 \bar{L} \bar{S} \} \times$$

$$\{ [\bar{J}] [L] [K] [S] \}^{\frac{1}{2}} \times$$

$$\left\{ \begin{matrix} \bar{S} & \bar{L} & \bar{J} \\ \ell & K & L \end{matrix} \right\} \left\{ \begin{matrix} L & \bar{S} & K \\ s & J & S \end{matrix} \right\} \mid \ell^4 \bar{L} \bar{S} \bar{J}, \ell; K, s, JM \rangle$$

where $\ell = 1$ and $L S J$ stands for the 2P_J ground state of iodine I. The symbol $(\ell^5 LS \{ \mid \ell^4 \bar{L} \bar{S} \})$ is the coefficient of fractional parentage³ and $[x] = 2x + 1$.

CONCLUSIONS:

The results, although approximate, are more or less in agreement with the experimentally observed laser lines⁴ with the exception of the 2056 cm^{-1} line which according to these calculations must have a strength much lower than that indicated by experiment. Experimentally, the $[^3\text{P}_0] \ 5d[2]_{5/2} \rightarrow [^3\text{P}_0] \ 6p[1]_{3/2}$ at 1818.61 cm^{-1} has a relative intensity of 5 and a transition probability of $0.23 \times 10^{+6} \text{ sec}^{-1}$. The $[^3\text{P}_2] \ 5d[4]_{7/2} \rightarrow [^3\text{P}_2] \ 6p[3]_{5/2}$ at 2915 cm^{-1} has a relative intensity of 10 and a transition probability of $0.91 \times 10^6 \text{ sec}^{-1}$ but the $[^3\text{P}_2] \ 5d[4]_{9/2} \rightarrow [^3\text{P}_2] \ 6p[3]_{7/2}$ at 2056 cm^{-1} has a relative intensity of 100 and a transition probability of $0.33 \times 10^6 \text{ sec}^{-1}$. However there are two other upper levels which have transitions at the same wavelength, namely

$$[3\text{P}_2] - 6f [3]_{7/2} \rightarrow [3\text{P}_2] - 7d [4]_{9/2}$$

and

$$[3\text{P}_2] - 9f [4]_{7/2} \rightarrow [3\text{P}_2] - 8d [4]_{7/2}$$

and the laser action could originate from these. This is somewhat unlikely since these two levels are lying far above the ground state. Other explanations such as cascade filling of the $[3\text{P}_2] - 5d [4]_{9/2}$ could be considered to explain the high intensity of the 2056 cm^{-1} transition.

TABLE 1a
TRANSITION PROBABILITIES FOR SOME OF THE
 $5p^4-5d \rightarrow 5p^4-6p$ STATES OF IODINE I

| ³ P] Core | | | | | | | |
|----------------------|-----|-----|----------------|---|---------|----------------------|------------------|
| Upper | | | Lower | | | A × 10 ⁻⁶ | wavelength |
| J _c | K | J | J _c | K | J | 1/sec | cm ⁻¹ |
| 0 | [2] | 3/2 | - | 0 | [1] 1/2 | 0.1939 | 1825.36 |
| | | | - | 0 | [1] 3/2 | 0.0231 | 1662.54 |
| 0 | [2] | 5/2 | - | 0 | [1] 3/2 | 0.2302 | 1818.61 |
| 1 | [1] | 1/2 | - | 1 | [0] 1/2 | 0.1640 | 1975.77 |
| | | | - | 1 | [1] 1/2 | 0.000008 | 90.09 |
| | | | - | 1 | [1] 3/2 | 0.0004 | 422.70 |
| | | | - | 1 | [2] 3/2 | 0.0003 | 670.11 |
| 1 | [1] | 3/2 | - | 1 | [0] 1/2 | 0.6248 | 3085.95 |
| | | | - | 1 | [1] 1/2 | 0.0046 | 1200.27 |
| | | | - | 1 | [1] 3/2 | 0.0478 | 1532.88 |
| | | | - | 1 | [2] 3/2 | 0.0006 | 1780.29 |
| | | | - | 1 | [2] 5/2 | 0.0083 | 2058.26 |
| 1 | [2] | 3/2 | - | 1 | [1] 1/2 | 0.3461 | 2436.78 |
| | | | - | 1 | [1] 3/2 | 0.1016 | 2769.39 |
| | | | - | 1 | [2] 3/2 | 0.2364 | 3016.80 |
| | | | - | 1 | [2] 5/2 | 0.0342 | 3294.77 |
| 1 | [2] | 5/2 | - | 1 | [1] 3/2 | 0.6605 | 2844.37 |
| | | | - | 1 | [2] 3/2 | 0.0188 | 3091.78 |
| | | | - | 1 | [2] 5/2 | 0.3417 | 3369.75 |
| 1 | [3] | 5/2 | - | 1 | [2] 3/2 | 0.4755 | 2370.11 |
| | | | - | 1 | [2] 5/2 | 0.0474 | 2648.08 |
| 1 | [3] | 7/2 | - | 1 | [2] 5/2 | 0.1163 | 1448.54 |
| 2 | [0] | 1/2 | - | 2 | [1] 1/2 | 0.2678 | 2758.79 |
| | | | - | 2 | [1] 3/2 | 0.0957 | 1553.66 |
| 2 | [1] | 1/2 | - | 2 | [1] 1/2 | 0.0573 | 1441.37 |
| | | | - | 2 | [1] 3/2 | 0.0001 | 236.24 |
| | | | - | 2 | [2] 3/2 | 0.1177 | 2308.33 |
| 2 | [1] | 3/2 | - | 2 | [1] 1/2 | 0.0006 | 498.14 |
| | | | - | 2 | [2] 3/2 | 0.0024 | 1365.10 |
| | | | - | 2 | [2] 5/2 | 0.0262 | 1448.79 |
| 2 | [2] | 3/2 | - | 2 | [1] 1/2 | 1.0157 | 4497.91 |
| | | | - | 2 | [1] 3/2 | 0.0797 | 3292.78 |
| | | | - | 2 | [2] 3/2 | 3.1022 | 5364.87 |
| | | | - | 2 | [2] 5/2 | 0.3611 | 5448.56 |
| | | | - | 2 | [3] 5/2 | 0.2666 | 4710.38 |

TABLE 1a (Cont.)

| Upper J _c K J | Lower J _c K J | A × 10 ⁻⁶ 1/sec | wavelength cm ⁻¹ |
|-----------------------------|-----------------------------|-------------------------------|--------------------------------|
| 2 [2] 5/2 | - 2 [1] 3/2 | 0.3948 | 3089.12 |
| | - 2 [2] 3/2 | 0.2046 | 5161.21 |
| | - 2 [2] 5/2 | 3.0060 | 5244.90 |
| | - 2 [3] 5/2 | 0.0111 | 4506.72 |
| | - 2 [3] 7/2 | 0.2186 | 4481.22 |
| 2 [3] 5/2 | - 2 [2] 3/2 | 0.0261 | 1030.48 |
| | - 2 [2] 5/2 | 0.0024 | 1114.17 |
| | - 2 [3] 5/2 | 0.0006 | 375.99 |
| | - 2 [3] 7/2 | 0.00003 | 350.49 |
| 2 [3] 7/2 | - 2 [2] 5/2 | 0.0348 | 1108.70 |
| | - 2 [3] 5/2 | 0.00002 | 370.52 |
| | - 2 [3] 7/2 | 0.0005 | 345.02 |
| 2 [4] 7/2 | - 2 [3] 5/2 | 0.9140 | 2915.02 |
| | - 2 [3] 7/2 | 0.0330 | 2889.52 |
| 2 [4] 9/2 | - 2 [3] 7/2 | 0.3328 | 2056.38 |

TABLE 1b
TRANSITION PROBABILITIES OF SOME OF THE
 $5p^4-6p \rightarrow 5p^4-6s$ STATES OF IODINE I

| [³ P] Core | | | | | | |
|-----------------------------|---|--|-----------------------------|-------------------------------|--------------------------------|--|
| Upper J _c K J | | | Lower J _c K J | A × 10 ⁻⁸ 1/sec | wavelength cm ⁻¹ | |
| 0 [1] 1/2 | - | | 0 [0] 1/2 | 0.4155 | 10917.67 | |
| 0 [1] 3/2 | - | | 0 [0] 1/2 | 0.4344 | 11080.49 | |
| 1 [0] 1/2 | - | | 1 [1] 1/2 | 0.0612 | 8314.72 | |
| | | | 1 [1] 3/2 | 0.1932 | 9681.71 | |
| 1 [1] 1/2 | - | | 1 [1] 1/2 | 0.2259 | 10200.40 | |
| | | | 1 [1] 3/2 | 0.1647 | 11567.39 | |
| 1 [1] 3/2 | - | | 1 [1] 1/2 | 0.0511 | 9867.79 | |
| | | | 1 [1] 3/2 | 0.3773 | 11234.78 | |
| 1 [2] 3/2 | - | | 1 [1] 1/2 | 0.2369 | 9626.38 | |
| | | | 1 [1] 3/2 | 0.0706 | 10987.37 | |
| 1 [2] 5/2 | - | | 1 [1] 3/2 | 0.3922 | 10709.40 | |
| 2 [1] 1/2 | - | | 2 [2] 3/2 | 0.2972 | 9764.08 | |
| 2 [1] 3/2 | - | | 2 [2] 3/2 | 0.0421 | 10969.21 | |
| | | | 2 [2] 5/2 | 0.5517 | 12428.63 | |
| 2 [2] 3/2 | - | | 2 [2] 3/2 | 0.2024 | 8897.12 | |
| | | | 2 [2] 5/2 | 0.0355 | 10356.54 | |
| 2 [2] 5/2 | - | | 2 [2] 3/2 | 0.0146 | 8813.43 | |
| | | | 2 [2] 5/2 | 0.3231 | 10272.85 | |
| 2 [3] 5/2 | - | | 2 [2] 3/2 | 0.2597 | 9551.61 | |
| | | | 2 [2] 5/2 | 0.0284 | 11011.03 | |
| 2 [3] 7/2 | - | | 2 [2] 5/2 | 0.4292 | 11036.53 | |

| [¹ D] core | | | | | | |
|------------------------|---|--|-----------|--------------------------|----------|--|
| | | | 2 [2] 5/2 | 0.3608 × 10 ⁸ | 10415.74 | |
| 2 [2] 5/2 | - | | 2 [2] 5/2 | 0.0213 × 10 ⁸ | 10004.68 | |
| | | | 2 [2] 3/2 | 0.2929 × 10 ⁸ | 9942.80 | |
| 2 [2] 5/2 | - | | 2 [2] 5/2 | 0.4577 × 10 ⁸ | 11537.58 | |
| | | | 2 [2] 3/2 | 0.0322 × 10 ⁸ | 11475.70 | |
| 2 [2] 3/2 | - | | 2 [2] 5/2 | 0.0480 × 10 ⁸ | 11451.95 | |
| | | | 2 [2] 3/2 | 0.4246 × 10 ⁸ | 11390.07 | |
| 2 [1] 3/2 | - | | 2 [2] 5/2 | 0.2727 × 10 ⁸ | 9827.38 | |
| | | | 2 [2] 3/2 | 0.0297 × 10 ⁸ | 9765.50 | |
| 2 [1] 1/2 | - | | 2 [2] 3/2 | 0.4310 × 10 ⁸ | 11051.84 | |

TABLE 1c
TRANSITION PROBABILITIES OF SOME OF THE
 $5p^4-6s-5p^5$ STATES OF IODINE I

| J_c | K | J | J_c | KJ | 3P Core | wavelength cm^{-1} |
|-------|-----|-----|-------|-------------|-----------------------------|-------------------------|
| | | | | | $A \times 10^{-8}$ 1/sec | |
| 0 | [0] | 1/2 | - | $^2P_{1/2}$ | 0.1117 | 53293.08 |
| | | | - | $^2P_{3/2}$ | 0.0833 | 60896.23 |
| 1 | [1] | 1/2 | - | $^2P_{1/2}$ | 0.4747 | 55583.60 |
| | | | - | $^2P_{3/2}$ | 0.4187 | 63186.75 |
| 1 | [1] | 3/2 | - | $^2P_{1/2}$ | 0.1323 | 54216.61 |
| | | | - | $^2P_{3/2}$ | 0.7346 | 61819.76 |
| 2 | [2] | 3/2 | - | $^2P_{1/2}$ | 0.0526 | 48489.73 |
| | | | - | $^2P_{3/2}$ | 0.2604 | 56092.88 |
| 2 | [2] | 5/2 | - | $^2P_{3/2}$ | 0.0902 | 54633.46 |

1D Core

| | | | | | | |
|---|-----|-----|---|-------------|--------|----------|
| 2 | [2] | 5/2 | - | $^2P_{3/2}$ | 0.3570 | 68587.87 |
| 2 | [2] | 3/2 | - | $^2P_{1/2}$ | 0.2098 | 61046.60 |
| | | | - | $^2P_{3/2}$ | 0.0597 | 68649.75 |

TABLE 1d
TRANSITION PROBABILITIES OF SOME OF THE
 $5p^4-5d \rightarrow 5p^5$ STATES OF IODINE I

| Upper $J_c K$ | J | Ground state | $A \times 10^{-8}$ 1/sec | wavelength cm^{-1} |
|------------------|-----|-----------------|-----------------------------|--------------------------------|
| 0 [2] | 1/2 | - $^2P_{1/2}$ | 0.7530 | 66036.11 |
| | | - $^2P_{3/2}$ | 0.0522 | 73637.26 |
| 0 [2] | 5/2 | - $^2P_{3/2}$ | 0.3153 | 73795.33 |
| 1 [1] | 1/2 | - $^2P_{1/2}$ | 1.6818 | 65874.09 |
| 1 [1] | 3/2 | - $^2P_{1/2}$ | 0.4421 | 66984.27 |
| | | - $^2P_{3/2}$ | 0.1221 | 74587.42 |
| 1 [2] | 3/2 | - $^2P_{1/2}$ | 0.4670 | 68220.78 |
| | | - $^2P_{3/2}$ | 0.1282 | 75823.93 |
| 1 [2] | 5/2 | - $^2P_{3/2}$ | 0.3430 | 75898.91 |
| 1 [3] | 5/2 | - $^2P_{3/2}$ | 1.1666 | 75177.24 |
| 2 [0] | 1/2 | - $^2P_{1/2}$ | 0.2970 | 61012.60 |
| | | - $^2P_{3/2}$ | 0.2112 | 68615.75 |
| 2 [1] | 1/2 | - $^2P_{1/2}$ | 0.4172 | 59695.18 |
| | | - $^2P_{3/2}$ | 1.1955 | 67298.33 |
| | | - $^2P_{1/2}$ | 0.0944 | 58751.95 |
| | | - $^2P_{3/2}$ | 0.0286 | 66355.10 |
| 2 [2] | 3/2 | - $^2P_{1/2}$ | 0.2827 | 62751.72 |
| | | - $^2P_{3/2}$ | 1.9920 | 70354.87 |
| 2 [3] | 5/2 | - $^2P_{3/2}$ | 1.5802 | 66020.48 |

TABLE 2
THE RADIAL INTEGRAL USED IN THE CALCULATIONS
OF THE TRANSITION PROBABILITIES IS
GIVEN BELOW IN ATOMIC UNITS

$$\int_0^{\infty} r f_{5d}(r) f_{6p}(r) dr = 46.44$$

$$\int_0^{\infty} r f_{6p}(r) f_{6s}(r) dr = 46.5$$

$$\int_0^{\infty} r f_{6s}(r) f_{5p}(r) dr = 0.8976$$

$$\int_0^{\infty} r f_{5d}(r) f_{5p}(r) dr = 1.69$$

TABLE 3
SOME OF THE RADIAL INTEGRALS OF $p^4-n\ell$ OF I.
FROM DEFINITIONS OF Yamanachi et al⁴ WE HAVE

$$G_{\ell-1} = 3G^{\ell-1}/2(2\ell+1)(2\ell-1)^2$$

$$G_{\ell+1} = 3G^{\ell+1}/2(2\ell+1)(2\ell+3)^2$$

and

$$F_2 = F^2/5(2\ell-1)(2\ell+1) .$$

$$F^2(5p^4 - 5d) = 14957.2 \text{ cm}^{-1}$$

$$G^1(5p^4 - 5d) = 14155.8 \text{ cm}^{-1}$$

$$G^3(5p^4 - 5d) = 8485.6 \text{ cm}^{-1}$$

$$\zeta_{5p} = 2678.4 \text{ cm}^{-1}$$

$$\zeta_{5d} = 46.4 \text{ cm}^{-1}$$

$$F^2(5p^4 - 6p) = 9474. \text{ cm}^{-1}$$

$$G^0(5p^4 - 6p) = 847 \text{ cm}^{-1}$$

$$G^2(5p^4 - 6p) = 1046 \text{ cm}^{-1}$$

$$\zeta_{5p} = 2739 \text{ cm}^{-1}$$

$$\zeta_{6p} = 124 \text{ cm}^{-1}$$

$$G^1(5p^4 - 6s) = 2292.9 \text{ cm}^{-1}$$

TABLE 4
LIFE TIMES OF SOME OF THE
EXCITED IODINE I LEVELS

$5p^4[{}^3P_{J_c}]5d[K] J$

| J_c | K | J | Energy level cm^{-1} | Lifetime (nsec) |
|-------|-----|-----|----------------------------------|--------------------|
| 0 | [2] | 3/2 | 73639.26 | 12.38 |
| 0 | [2] | 5/2 | 73795.33 | 31.49 |
| 1 | [1] | 1/2 | 73477.24 | 5.94 |
| 1 | [1] | 3/2 | 74587.42 | 17.52 |
| 1 | [2] | 3/2 | 75823.93 | 16.60 |
| 1 | [2] | 5/2 | 75898.91 | 28.31 |
| 1 | [3] | 5/2 | 75177.24 | 8.53 |
| 1 | [3] | 7/2 | 73977.70 | 8620.7 |
| 2 | [0] | 1/2 | 68615.75 | 19.54 |
| 2 | [1] | 1/2 | 67298.38 | 6.19 |
| 2 | [1] | 3/2 | 66355.10 | 77.94 |
| 2 | [2] | 3/2 | 70354.87 | 4.30 |
| 2 | [2] | 5/2 | 70151.21 | 261.1 |
| 2 | [3] | 5/2 | 66020.48 | 6.33 |
| 2 | [3] | 7/2 | 66015.01 | 28735.6 |
| 2 | [4] | 7/2 | 68559.51 | 1063.8 |
| 2 | [4] | 9/2 | 67726.37 | 3030.3 |

$5p^4-[]{}^3P_{J_c}]6p[K] J$

| | | | | |
|---|-----|-----|----------|------|
| 0 | [1] | 1/2 | 71813.90 | 24.1 |
| 0 | [1] | 3/2 | 71976.72 | 23.0 |
| 1 | [0] | 1/2 | 71501.47 | 39.4 |
| 1 | [1] | 1/2 | 73387.15 | 25.6 |
| 1 | [1] | 3/2 | 73054.54 | 23.3 |
| 1 | [2] | 3/2 | 72807.13 | 32.6 |
| 1 | [2] | 5/2 | 72529.16 | 25.5 |
| 2 | [1] | 1/2 | 65856.96 | 33.7 |
| 2 | [1] | 3/2 | 67062.09 | 16.7 |
| 2 | [2] | 3/2 | 64990.00 | 42.1 |
| 2 | [2] | 5/2 | 64906.31 | 29.6 |
| 2 | [3] | 5/2 | 65644.49 | 34.7 |
| 2 | [3] | 7/2 | 65669.99 | 23.3 |

TABLE 4 (Cont.)

 $5p^4 [^3P_{J_C}] 6s [K] J$

| J_C | K | J | Energy level cm^{-1} | Lifetimes (nsec) |
|-------|-----|-----|----------------------------------|---------------------|
| 0 | [0] | 1/2 | 60896.23 | 51.3 |
| 1 | [1] | 1/2 | 63186.75 | 23.9 |
| 1 | [1] | 3/2 | 61819.76 | 75.6 |
| 2 | [2] | 3/2 | 56092.88 | 32.0 |
| 2 | [2] | 5/2 | 54633.46 | 111.0 |

 $5p^4 - [^1D_2] - 6p [K] J$

| | | | |
|-----|-----|----------|------|
| [1] | 1/2 | 79701.59 | 23.2 |
| [1] | 3/2 | 78415.25 | 33.1 |
| [2] | 3/2 | 80039.82 | 21.2 |
| [2] | 5/2 | 80125.45 | 20.2 |
| [3] | 5/2 | 78592.55 | 31.8 |
| [3] | 7/2 | 77003.61 | 27.7 |

 $5p^4 [^1D_2] 6s [K] J$

| | | | |
|-----|-----|----------|------|
| [2] | 3/2 | 68549.75 | 37.2 |
| [2] | 5/2 | 68587.87 | 28.0 |

REFERENCES

1. Minnahan, L., Arkiv. For Fysik, Band 21, No. 26.
2. Herman, F. and S. Skilman, Atomic Structure Calculations, Englewood Cliff, N.J., (1963).
3. Nielsen, C.W. and G.F. Koster, Spectroscopic Coefficients for the p^n , d^n and f^n Configuration, The M.I.T. Press, (1963).
4. Yamanachi, J. and A. Amemiya, Proceedings of Physical Society of Japan, Vol. I, page 18, (1946).